

**TEAT GUARD SANITIZING TEAT DIP - iodine solution**  
Ecolab Inc.

**FOR COMMERCIAL USE ONLY**  
**KEEP OUT OF REACH OF CHILDREN**  
**NOT FOR HUMAN USE**

**CAUTION**

Not for internal use. May cause eye irritation. If swallowed, may cause irritation, nausea, stomach distress.

**PRECAUTIONARY STATEMENTS**

No protective equipment is necessary under normal use conditions. If splashing may occur, eye protection is recommended.

**FIRST AID**

**If in Eyes:** Immediately flush eyes with plenty of cool running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.

**If Swallowed:** Rinse mouth; then immediately drink 1 or 2 large glasses of water. DO NOT induce vomiting. Never give anything by mouth to an unconscious person. IF IRRITATION OR DISCOMFORT PERSISTS, CALL A PHYSICIAN.

**FOR EMERGENCY MEDICAL INFORMATION IN USA OR CANADA, CALL: 1-800-328-0026.**

**FOR EMERGENCY MEDICAL INFORMATION WORLDWIDE, CALL: 1-651-225-3225 (IN THE USA).**

**DIRECTIONS FOR USE**

**IMPORTANT:** Do not mix *Teat Guard* with water, any other teat dips, or other products unless specified in product use instructions. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for *Teat Guard*.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

**Before Milking:** Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip or teat spray to avoid contamination of milk. Teats should then be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.

**Pre-Milk Dipping:** Fill teat dip cup with *Teat Guard*. Do not dilute. Before each cow is milked, dip each teat full-length into the teat dip cup containing *Teat Guard*. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

**Post-Milk Dipping:** Immediately after each milking, use *Teat Guard* at full-strength. Dip each teat full-length into the teat dip cup containing *Teat Guard*. Allow to air dry. Do not wipe. Always use fresh, full-strength *Teat Guard*. Do not turn cows out in freezing weather until *Teat Guard* is completely dry.

**Expanded Usage:** When freshening cows, begin dipping teats twice daily for about 10 days before calving.

**PRECAUTION:** *Teat Guard* is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

**PRINCIPLE DISPLAY PANEL/REPRESENTATIVE LABEL**

**ECOLAB**

**42325**

**Net Contents: 15 U.S. gal/56.78 L**

**Teat Guard**

**Sanitizing Teat Dip**

An Aid in Reducing the Spread of Organisms Which May Cause Mastitis

**ACTIVE INGREDIENT:**

Iodine ..... 1.0%

**INACTIVE INGREDIENTS:** ..... 99.0%

(Contains an emollient system.)

**INDICATIONS, DOSAGE AND ADMINISTRATION:**

As an aid in the prevention or control of losses due to CRD associated with *M. gallisepticum* (PPL0). Two (2) g/gal spectinomycin activity in the drinking water for the first three days of life and for one day following each vaccination.

As an aid in controlling infectious synovitis due to *Mycoplasma synoviae* in broiler chickens. One (1) g/gal spectinomycin activity in the drinking water for the first three to five days of life.

For increased rate of weight gain and improved feed efficiency in floor-raised broiler chickens. One-half (1/2) g/gal spectinomycin activity for the first three days of life and for one day following each vaccination.

**DIRECTIONS FOR USE:**

To provide 2 g/gal: Dissolve contents in 250 gallons of water. For proportioners, dissolve contents of one-half (1/2) package in one (1) gallon of water. Meter at one (1) oz/gal.

To provide 1 g/gal: Dissolve contents in 500 gallons of water. For proportioners, dissolve contents of one-half (1/2) package in one (2) gallons of water. Meter at one (1) oz/gal.

To provide 1/2 g/gal: Dissolve contents of one-half (1/2) package in 500 gallons of water. For proportioners, dissolve contents of one-half (1/2) package in one (4) gallons of water. Meter at one (1) oz/gal.

# Spectam®

(Spectinomycin)

WATER SOLUBLE CONCENTRATE

ANTI-INFECTIVE

for growing chickens

Each 1000 g contains: Spectinomycin dihydrochloride pentahydrate equivalent to spectinomycin activity ..... 500.00 g

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

NADA 038-361, Approved by FDA

**NET CONTENTS:**  
**1000 g (35.27 oz)**

Manufactured by:  
Bimeda, Inc.  
Le Sueur, MN 56058

**Osborn®**

**WARNING:** DO NOT administer this drug within 5 days of slaughter. DO NOT administer to laying chickens.


**PRECAUTION:** Rarely, some individuals who handle spectinomycin develop serious reactions involving skin, nails and eyes. Individuals who have experienced rash or other evidence of allergic reaction should avoid further contact with spectinomycin.

Store at 20°C - 25°C (68°F - 77°F), excursions permitted between 15°C - 30°C (59°F - 86°F).



© Registered Trademark of Bimeda, Inc.  
Bimeda, Inc. is a division of Cross Veipharma Group Ltd.

8SPE001-809 Product No. 13PE001



LOT:  
EXP:

